

Total number of printed pages-23

3 (Sem-5/CBCS) MAT HC 1 (N/O)

2022

**MATHEMATICS**

(Honours)

Paper : MAT-HC-5016

(For New Syllabus)

(Complex Analysis)

Full Marks : 60

Time : Three hours

***The figures in the margin indicate full marks for the questions.***

1. Answer **any seven** questions from the following : 1×7=7

(a) Describe the domain of definition of the

function  $f(z) = \frac{z}{z + \bar{z}}$ .

(b) What is the multiplicative inverse of a non-zero complex number  $z = (x, y)$  ?

Contd.

- (c) Verify that  $(3, 1) (3, -1) \left(\frac{1}{5}, \frac{1}{10}\right) = (2, 1)$ .
- (d) Determine the accumulation points of the set  $Z_n = \frac{i}{n} (n = 1, 2, 3, \dots)$ .
- (e) Write the Cauchy-Riemann equations for a function  $f(z) = u + iv$ .
- (f) When a function  $f$  is said to be analytic at a point?
- (g) Determine the singular points of the function  $f(z) = \frac{2z + 1}{z(z^2 + 1)}$ .
- (h)  $\exp(2 \pm 3\pi i)$  is
- (i)  $-e^2$
  - (ii)  $e^2$
  - (iii)  $2e$
  - (iv)  $-2e$  (Choose the correct answer)

(i) The value of  $\log(-1)$  is

(i) 0

(ii)  $2n\pi i$

(iii)  $\pi i$

(iv)  $-\pi i$  (Choose the correct answer)

(j) If  $z = x + iy$ , then  $\sin z$  is

(i)  $\sin x \cos hy + i \cos x \sin hy$

(ii)  $\cos x \cos hy - i \sin x \sin hy$

(iii)  $\cos x \sin hy + i \sin x \cos hy$

(iv)  $\sin x \sin hy - i \cos x \cos hy$

(Choose the correct answer)

(k) If  $\cos z = 0$ , then

(i)  $z = n\pi, (n = 0, \pm 1, \pm 2, \dots)$

(ii)  $z = \frac{\pi}{2} + n\pi, (n = 0, \pm 1, \pm 2, \dots)$

(iii)  $z = 2n\pi, (n = 0, \pm 1, \pm 2, \dots)$

(iv)  $z = \frac{\pi}{2} + 2n\pi, (n = 0, \pm 1, \pm 2, \dots)$

(Choose the correct answer)

(l) If  $z_0$  is a point in the  $z$ -plane, then

$$\lim_{z \rightarrow \infty} f(z) = \infty \text{ if}$$

$$(i) \quad \lim_{z \rightarrow 0} \frac{1}{f(z)} = \infty$$

$$(ii) \quad \lim_{z \rightarrow 0} f\left(\frac{1}{z}\right) = 0$$

$$(iii) \quad \lim_{z \rightarrow 0} \frac{1}{f(z)} = 0$$

$$(iv) \quad \lim_{z \rightarrow 0} \frac{1}{f\left(\frac{1}{z}\right)} = 0$$

(Choose the correct answer)

2. Answer **any four** questions from the following : 2×4=8

(a) Reduce the quantity  $\frac{5i}{(1-i)(2-i)(3-i)}$

to a real number.

(b) Define a connected set and give *one* example.

(c) Find all values of  $z$  such that  $\exp(2z-1)=1$ .

(d) Show that  $\log(i^3) \neq 3\log i$ .

(e) Show that

$$2\sin(z_1+z_2)\sin(z_1-z_2) = \cos 2z_2 - \cos 2z_1$$

(f) If  $z_0$  and  $w_0$  are points in the  $z$  plane and  $w$  plane respectively, then prove that  $\lim_{z \rightarrow z_0} f(z) = \infty$  if and only if

$$\lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0.$$

(g) State the Cauchy integral formula. Find

$$\frac{1}{2\pi i} \int_C \frac{1}{z-z_0} dz \text{ if } z_0 \text{ is any point}$$

interior to simple closed contour  $C$ .

(h) Show that  $\int_0^{\frac{\pi}{6}} e^{i2t} dt = \frac{\sqrt{3}}{4} + \frac{i}{4}$ .

3. Answer **any three** questions from the following : 5×3=15

- (a) (i) If  $a$  and  $b$  are complex constants, use definition of limit to show that

$$\lim_{z \rightarrow z_0} (az + b) = az_0 + b. \quad 2$$

- (ii) Show that

$$\lim_{z \rightarrow 0} \left( \frac{z}{\bar{z}} \right)^2 \text{ does not exist.} \quad 3$$

- (b) Suppose that  $\lim_{z \rightarrow z_0} f(z) = w_0$  and

$$\lim_{z \rightarrow z_0} F(z) = W_0.$$

Prove that  $\lim_{z \rightarrow z_0} [f(z)F(z)] = w_0W_0$ .

- (c) (i) Show that for the function  $f(z) = \bar{z}$ ,  $f'(z)$  does not exist anywhere. 3

(ii) Show that  $\lim_{z \rightarrow \infty} \frac{4z^2}{(z-1)^2} = 4$ . 2

(d) (i) Show that the function  
 $f(z) = \exp \bar{z}$  is not analytic  
anywhere. 3

(ii) Find all roots of the equation  
 $\log z = i \frac{\pi}{2}$ . 2

(e) If a function  $f$  is analytic at all  
points interior to and on a simple  
closed contour  $C$ , then prove that

$$\int_C f(z) dz = 0.$$

(f) Evaluate :  $2^{1/2} + 2^{1/2} = 5$

(i)  $\int_C \frac{e^{-z}}{z - (\pi i/2)} dz$

(ii)  $\int_C \frac{z}{2z + 1} dz$

where  $C$  denotes the positively oriented  
boundary of the square whose sides lie  
along the lines  $x = \pm 2$  and  $y = \pm 2$ .

(g) Prove that any polynomial

$$P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n \quad (a_n \neq 0).$$

of degree  $n$  ( $n \geq 1$ ) has at least one zero.

(h) Find the Laurent series that represents

$$\text{the function } f(z) = z^2 \sin\left(\frac{1}{z^2}\right) \text{ in the}$$

domain  $0 < |z| < \infty$ .

4. Answer **any three** questions from the following : 10×3=30

(a) (i) If a function  $f$  is continuous throughout a region  $R$  that is both closed and bounded, then prove that there exists a non-negative real number  $\mu$  such that  $|f(z)| \leq \mu$  for all points  $z$  in  $R$ , where equality holds for at least one such  $z$ .

4



(ii) Let a function  $f(z) = u(x, y) + iv(x, y)$  be analytic throughout a given domain  $D$ . If  $|f(z)|$  is constant throughout  $D$ , then prove that  $f(z)$  must be constant there too. 3

(iii) Show that the function  $f(z) = \sin x \cosh y + i \cos x \sinh y$  is entire. 3

(b) (i) Suppose that  $f(z_0) = g(z_0) = 0$  and that  $f'(z_0)$   $g'(z_0)$  exist, where  $g'(z_0) \neq 0$ . Use definition of derivative to show that

$$\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = \frac{f'(z_0)}{g'(z_0)}. \quad 3$$

(ii) Show that  $f'(z)$  does not exist at any point if  $f(z) = 2x + ixy^2$ . 3

(iii) If a function  $f$  is analytic at a given point, then prove that its derivatives of all orders are analytic there too. 4

(c) Let the function

$f(z) = u(x, y) + iv(x, y)$  be defined throughout some  $\varepsilon$ -neighbourhood of a point  $z_0 = x_0 + iy_0$ . If  $u_x, u_y, v_x, v_y$  exist everywhere in the neighbourhood, and these partial derivatives are continuous at  $(x_0, y_0)$  and satisfy the Cauchy-Riemann equations at  $(x_0, y_0)$ , then prove that  $f'(z_0)$  exist and  $f'(z_0) = u_x + iv_x$  where the right hand side is to be evaluated at  $(x_0, y_0)$ .

Use it to show that for the function

$f(z) = e^{-x} \cdot e^{-y}$ ,  $f''(z)$  exists everywhere and  $f''(z) = f(z)$ . 6+4=10

(d) (i) Prove that the existence of the derivative of a function at a point implies the continuity of the function at that point.

With the help of an example show that the continuity of a function at a point does not imply the existence of derivative there.

3+5=8

(ii) Find  $f'(z)$  if

$$f(z) = \frac{z-1}{2z+1} \left( z \neq -\frac{1}{2} \right). \quad 2$$

(e) (i) Prove that  $\int_C \frac{dz}{z} = \pi i$  where  $C$  is

the right-hand half  $z = 2e^{i\theta}$

$\left( -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2} \right)$  of the circle  $|z| = 2$

from  $z = -2i$  to  $z = 2i$ . 5

(ii) If a function  $f$  is analytic everywhere inside and on a simple closed contour  $C$ , taken in the positive sense, then prove that

$$f'(z) = \frac{1}{2\pi i} \int_C \frac{f(s)}{(s-z)^2} ds \quad \text{where } s$$

denotes points on  $C$  and  $z$  is interior to  $C$ . 5

(f) (i) Evaluate  $I = \int_C z^{a-1} dz$

where  $C$  is the positively oriented circle  $z = Re^{i\theta}$  ( $-\pi \leq \theta \leq \pi$ ) about the origin and  $a$  denote any non-zero real number.

If  $a$  is a non-zero integer  $n$ , then what is the value of  $\int_C z^{n-1} dz$  ?

$$4+1=5$$

(ii) Let  $C$  denote a contour of length  $L$ , and suppose that a function  $f(z)$  is piecewise continuous on  $C$ . If  $\mu$  is a non-negative constant such that  $|f(z)| \leq \mu$  for all point  $z$  on  $C$  at which  $f(z)$  is defined, then prove

$$\text{that } \left| \int_C f(z) dz \right| \leq \mu L.$$

Use it to show that  $\left| \int_C \frac{dz}{z^2 - 1} \right| \leq \frac{\pi}{3}$

where  $C$  is the arc of the circle  $|z| = 2$  from  $z = 2$  to  $z = 2i$  that lies in the 1st quadrant.

$$3+2=5$$

(g) (i) Apply the Cauchy-Goursat theorem to show that  $\int_C f(z) = 0$  when the contour  $C$  is the unit circle  $|z|=1$ , in either direction and  $f(z) = ze^{-z}$ . 4

(ii) If  $C$  is the positively oriented unit circle  $|z|=1$  and  $f(z) = \exp(2z)$  find  $\int_C \frac{f(z)}{z^4} dz$ . 3

(iii) Let  $z_0$  be any point interior to a positively oriented simple closed curve  $C$ . Show that

$$\int_C \frac{dz}{(z - z_0)^{n+1}} = 0, (n = 1, 2, \dots). \quad 3$$

(h) (i) Suppose that  $z_n = x_n + iy_n$ ,  $(n = 1, 2, \dots)$  and  $z = x + iy$ . Prove that  $\lim_{n \rightarrow \infty} z_n = z$  if and only if

$$\lim_{n \rightarrow \infty} x_n = x \quad \text{and} \quad \lim_{n \rightarrow \infty} y_n = y. \quad 5$$

(ii) Show that

$$z^2 e^{3z} = \sum_{n=2}^{\infty} \frac{3^{n-2}}{(n-2)!} z^n \quad (|z| < \infty)$$

5

( For Old Syllabus )

(Riemann Integration and Metric Spaces)

Full Marks : 80

Time : Three hours

**The figures in the margin indicate full marks for the questions.**

1. Answer the following questions :  $1 \times 10 = 10$

(a) Describe an open ball on the real line  $\mathbb{R}$  for the usual metric  $d$ .

(b) Find the limit point of the set

$$\left\{ 1, \frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{n}, \dots \right\}.$$

(c) Define Cauchy sequence in a metric space  $(X, d)$ .

(d) Let  $A$  and  $B$  be two subsets of a metric space  $(X, d)$ . Then

$$(i) \quad (A \cap B)^0 = A^0 \cap B^0$$

$$(ii) \quad (A \cup B)^0 = A^0 \cup B^0$$

$$(iii) \quad (A \cap B)' = A' \cap B'$$

$$(iv) \quad (A \cup B)' = A' \cup B'$$

where  $A^0$  denotes interior of  $A$

$A'$  denotes derived set of  $A$

(Choose the correct answer)

- (e) In a complete metric space
- (i) every sequence is bounded
  - (ii) every bounded sequence is convergent
  - (iii) every convergent sequence is bounded
  - (iv) every Cauchy sequence is convergent
- (Choose the correct answer)*

(f) Let  $\{F_n\}$  be a decreasing sequence of closed subsets of a complete metric space and  $d(F_n) \rightarrow 0$  as  $n \rightarrow \infty$ . Then

(i)  $\bigcap_{n=1}^{\infty} F_n = \phi$

(ii)  $\bigcap_{n=1}^{\infty} F_n$  contains at least one point

(iii)  $\bigcap_{n=1}^{\infty} F_n$  contains exactly one point

(iv)  $d\left(\bigcap_{n=1}^{\infty} F_n\right) > 0$

*(Choose the correct answer)*

(g) Let  $(X, d)$  and  $(Y, \rho)$  be metric spaces and  $A \subset X$ . Let  $f : X \rightarrow Y$  be continuous on  $X$ . Then

(i)  $f(A) = f(\overline{A})$

(ii)  $f(\overline{A}) \subset \overline{f(A)}$

(iii)  $\overline{f(A)} \subset f(\overline{A})$

(iv)  $f(A) = f(A^0)$

*(Choose the correct answer)*

(h) What is meant by partition  $P$  of an interval  $[a, b]$ ?

(i) Prove that  $\overline{\alpha + 1} = \alpha \overline{\alpha}$

(j) Define the upper and the lower Darboux sums of a function  $f : [a, b] \rightarrow \mathbb{R}$  with respect to a partition  $P$ .

2. Answer the following questions :  $2 \times 5 = 10$

(a) Prove that in a discrete metric space every singleton set is open.



- (b) For any two subsets  $F_1$  and  $F_2$  of a metric space  $(X, d)$ , prove that

$$\overline{(F_1 \cup F_2)} = \overline{F_1} \cup \overline{F_2}$$

- (c) Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces and let  $f : X \rightarrow Y$ . Then if  $f$  is continuous on  $X$ , prove that

$$\overline{f^{-1}(B)} \subset f^{-1}(\overline{B}) \text{ for all subsets } B \text{ of } Y.$$

- (d) Find  $L(f, P)$  and  $U(f, P)$  for a constant function  $f : [a, b] \rightarrow \mathbb{R}$ .

- (e) Examine the existence of improper

$$\text{integral } \int_0^1 \frac{1}{\sqrt{x}} dx.$$

3. Answer **any four** parts : 5×4=20

- (a) Let  $d$  be a metric on the non-empty set  $X$ . Prove that the function  $d'$  defined by

$$d'(x, y) = \min\{1, d(x, y)\}$$

where  $x, y \in X$  is a metric on  $X$ . State whether  $d'$  is bounded or not.

$$4+1=5$$

- (b) In a metric space  $(X, d)$ , prove that every closed sphere is a closed set.
- (c) Prove that if a Cauchy sequence of points in a metric space  $(X, d)$  contains a convergent subsequence, then the sequence also converges to the same limit as the subsequence.
- (d) Let  $(X, d)$  be a metric space and let  $\{Y_\lambda, \lambda \in I\}$  be a family of connected sets in  $(X, d)$  having a non-empty intersection. Then prove that  $Y = \bigcup_{\lambda \in I} Y_\lambda$  is connected.
- (e) Consider the function  $f : [0, 1] \rightarrow \mathbb{R}$  defined by  $f(x) = \begin{cases} 1 & \text{if } x \in Q \\ 0 & \text{otherwise} \end{cases}$   
Prove that  $f$  is not integrable on  $[0, 1]$ .
- (f) Let  $f : [a, b] \rightarrow \mathbb{R}$  be bounded and monotone. Prove that  $f$  is integrable.

4. Answer **any four** parts :  $10 \times 4 = 40$

(a) (i) Define a metric space.

Let

$$X = \mathbb{R}^n = \{x = (x_1, x_2, \dots, x_n), x_i \in \mathbb{R}, 1 \leq i \leq n\}$$

be the set of all real  $n$ -tuples.

For  $x = (x_1, x_2, \dots, x_n)$  and

$y = (y_1, y_2, \dots, y_n)$  in  $\mathbb{R}^n$  define

$$d(x, y) = \left( \sum_{i=1}^n (x_i - y_i)^2 \right)^{1/2}.$$

Prove that  $(\mathbb{R}^n, d)$  is a metric space.  $2+4=6$

(ii) Prove that in a metric space  $(X, d)$ , a finite intersection of open sets is open.  $4$

(b) Let  $Y$  be a subspace of a metric space  $(X, d)$ . Prove the following :  $5+5=10$

(i) Every subset of  $Y$  that is open in  $Y$  is also open in  $X$  if and only if  $Y$  is open in  $X$ .

(ii) Every subset of  $Y$  that is closed in  $Y$  is also closed in  $X$  if and only if  $Y$  is closed in  $X$ .

(c) (i) Prove that the function

$f : [0, 1] \rightarrow \mathbb{R}$  defined by

$f(x) = x^2$  is uniformly

continuous. Further prove that the function will not be uniformly continuous if the domain is  $\mathbb{R}$ .

3+3=6

(ii) Let  $(X, d_X)$ ,  $(Y, d_Y)$  and  $(Z, d_Z)$  be metric spaces and let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  be continuous. Prove that the composition  $g \circ f$  is a continuous map of  $X$  into  $Z$ .

4

(d) When a metric space is said to be disconnected ?

Prove that a metric space  $(X, d)$  is disconnected if and only if there exists a non-empty proper subset of  $X$  which is both open and closed in  $(X, d)$ .

2+8=10

- (e) (i) Show that the metric space  $(X, d)$  where  $X$  denotes the space of all sequences  $x = (x_1, x_2, x_3, \dots, x_n)$  of real numbers for which

$$\left( \sum_{k=1}^{\infty} |x_k|^p \right)^{\frac{1}{p}} < \infty \quad (p \geq 1) \text{ and } d \text{ is the}$$

metric given by

$$d_p(x, y) = \left( \sum_{k=1}^{\infty} (x_k - y_k)^p \right)^{1/p}, \quad x, y \in X$$

is a complete metric space. 7

- (ii) Let  $X$  be any non-empty set and let  $d$  be defined by

$$d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

Show that  $(X, d)$  is a complete metric space. 3

(f) Prove that a bounded function  $f : [a, b] \rightarrow \mathbb{R}$  is integrable if and only if for each  $\varepsilon > 0$ , there exists a partition  $P$  of  $[a, b]$  such that  $U(P, f) - L(P, f) < \varepsilon$ .

(g) Let  $f : [0, 1] \rightarrow \mathbb{R}$  be continuous. Let

$C_i \in \left[ \frac{i-1}{n}, \frac{i}{n} \right], n \in \mathbb{N}$ . Then prove that

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(C_i) = \int_0^1 f(x) dx.$$

Using it, prove that  $\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{n}{k^2 + n^2} = \frac{\pi}{4}$ .

$$6+4=10$$

(h) (i) Prove that a mapping  $f : X \rightarrow Y$  is continuous on  $X$  if and only if  $f^{-1}(F)$  is closed in  $X$  for all closed subsets  $F$  of  $Y$ . 5

(ii) Let  $f$  and  $g$  be continuous on  $[a, b]$ . Also assume that  $g$  does not change sign on  $[a, b]$ . Then prove that for some  $c \in [a, b]$  we have

$$\int_a^b f(x) g(x) dx = f(c) \int_a^b g(x) dx.$$

5

