

Total number of printed pages-16

3 (Sem-5/CBCS) MAT HC 1 (N/O)

2023

MATHEMATICS

(Honours Core)

OPTION-A

(For New Syllabus)

Paper : MAT-HC-5016

(Complex Analysis)

Full Marks : 60

Time : Three hours

The figures in the margin indicate full marks for the questions.

1. Answer the following questions : $1 \times 7 = 7$

(a) Which point on the Riemann sphere represents ∞ of the extended complex plane $\mathbb{C} \cup \{\infty\}$?

(b) A set $S \subseteq \mathbb{C}$ is closed if and only if S contains each of its _____ points.

(Fill in the gap)

Contd.

- (c) Write down the polar form of the Cauchy-Riemann equations.
- (d) The function $f(z) = \sinh z$ is a periodic function with a period _____ .
(Fill in the gap)
- (e) Define a simple closed curve.
- (f) Write down the value of the integral $\int_C f(z) dz$, where $f(z) = ze^{-2}$ and C is the circle $|z| = 1$.
- (g) Find $\lim_{n \rightarrow \infty} z_n$, where $z_n = -1 + i \frac{(-1)^n}{n^2}$.

2. Answer the following questions : $2 \times 4 = 8$

- (a) Let $f(z) = i \frac{z}{2}$, $|z| < 1$. Show that

$$\lim_{z \rightarrow 1} f(z) = \frac{i}{2}, \text{ using } \varepsilon - \delta \text{ definition.}$$

- (b) Show that all the zeros of $\sinh z$ in the complex plane lie on the imaginary axis.

(c) Evaluate the contour integral

$$\int_C \frac{dz}{z}, \text{ where } C \text{ is the semi circle}$$
$$z = e^{i\theta}, \quad 0 \leq \theta \leq \pi$$

(d) Using Cauchy's integral formula, evaluate

$$\int_C \frac{e^{2z}}{z^4} dz, \text{ where } C \text{ is the circle } |z| = 1.$$

3. Answer **any three** questions from the following : 5×3=15

(a) Find all the fourth roots of -16 and show that they lie at the vertices of a square inscribed in a circle centered at the origin.

(b) Suppose $f(z) = u(x, y) + iv(x, y)$,
($z = x + iy$) and $z_0 = x_0 + iy_0$,
 $w_0 = u_0 + iv_0$. Then prove the following:

$$\lim_{(x, y) \rightarrow (x_0, y_0)} u(x, y) = u_0,$$

$$\lim_{(x, y) \rightarrow (x_0, y_0)} v(x, y) = v_0, \text{ if and only}$$

$$\text{if } \lim_{z \rightarrow z_0} f(z) = w_0.$$

(c) (i) Show that the function $f(z) = \operatorname{Re} z$ is nowhere differentiable.

(ii) Let $T(z) = \frac{az+b}{cz+d}$, where

$$ad - bc \neq 0.$$

Show that $\lim_{z \rightarrow \infty} T(z) = \infty$ if $c = 0$.

$$3+2=5$$

(d) Let C be the arc of the circle $|z|=2$ from $z=2$ to $z=2i$ that lies in the first quadrant. Show that

$$\left| \int_C \frac{z+4}{z^3-1} dz \right| \leq \frac{6\pi}{7}$$

(e) State and prove fundamental theorem of algebra.

4. Answer **any three** questions from the following : 10×3=30

(a) (i) Show that $\exp(z + \pi i) = -\exp(z)$

1

(ii) Show that

$$\log(-1+i)^2 \neq 2\log(-1+i) \quad 2$$

(iii) Show that

$$|\sin z|^2 = \sin^2 x + \sinh^2 y \quad 2$$

(iv) Show that a set $S \subseteq \mathbb{C}$ is unbounded if and only if every neighbourhood of the point at infinity contains at least one point of S . 5

(b) (i) Suppose that $f(z_0) = g(z_0) = 0$ and that $f'(z_0)$, $g'(z_0)$ exist with $g'(z_0) \neq 0$. Using the definition of derivative show that

$$\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = \frac{f'(z_0)}{g'(z_0)} \quad 5$$

(ii) Show that

$$z^2 e^{3z} = \sum_{n=2}^{\infty} \frac{3^{n-2}}{(n-2)!} z^n,$$

where $|z| < \infty$. 5

(c) State and prove Laurent's theorem.

(d) (i) Using definition of derivative, show that $f(z) = |z|^2$ is nowhere differentiable except at $z = 0$. 5

- (ii) Define singular points of a function. Determine singular points of the functions :

$$f(z) = \frac{2z+1}{z(z^2+1)} ;$$

$$g(z) = \frac{z^3+i}{z^2-3z+2} \quad 1+4=5$$

- (e) (i) Let $f(z) = u(x, y) + iv(x, y)$ be analytic in a domain D . Prove that the families of curves $u(x, y) = c_1$, $v(x, y) = c_2$ are orthogonal.

- (ii) Let C denote a contour of length L and suppose that a function $f(z)$ is piecewise continuous on C . If M is a non-negative constant such that

$$|f(z)| \leq M \text{ for all } z \text{ in } C$$

then show that

$$\left| \int_C f(z) dz \right| \leq ML. \quad 5+5=10$$

- (f) (i) Prove that two non-zero complex numbers z_1 and z_2 have the same moduli if and only if $z_1 = c_1 c_2$, $z_2 = c_1 \bar{c}_2$, for some complex numbers c_1, c_2 . 4
- (ii) Show that mean value theorem of integral calculus of real analysis does not hold for complex valued functions $w(t)$. 3
- (iii) State Cauchy-Goursat theorem. 1
- (iv) Show that $\lim_{z \rightarrow \infty} \frac{z^2 + 1}{z - 1} = \infty$. 2
-

OPTION-B

(For Old Syllabus)

(Riemann Integration and Metric Spaces)

Full Marks : 80

Time : Three hours

The figures in the margin indicate full marks for the questions.

1. Answer the following questions : $1 \times 10 = 10$
 - (a) Write the statement of the First Fundamental Theorem of Calculus.
 - (b) Evaluate $\int_0^{\infty} e^{-x} dx$, if it exists.
 - (c) Prove that $\Gamma(1) = 1$.
 - (d) Define a complete metric space.
 - (e) Describe an open ball in the discrete metric space (X, d) .
 - (f) $(A \cup B)^0$ need not be $A^0 \cup B^0$ — Justify it where A and B are subsets of a metric space (X, d) .
 - (g) Find the derived sets of the intervals $(0, 1)$ and $[0, 1]$.

(h) Let A and B be two subsets of a metric space (X, d) . Which of the following is not correct?

(i) $A \subseteq B \Rightarrow A' \subseteq B'$

(ii) $(A \cap B)' \subseteq A' \cap B'$

(iii) $A' \cap B' \subseteq (A \cap B)'$

(iv) $(A \cup B)' = A' \cup B'$

(i) The Euclidean metric on \mathbb{R}^n is defined as

(i) $d(x, y) = \left\{ \sum_{i=1}^n (x_i - y_i)^2 \right\}^{\frac{1}{2}}$

(ii) $d(x, y) = \left\{ \sum_{i=1}^n |x_i - y_i|^p \right\}^{\frac{1}{p}}$

where $p \geq 1$

(iii) $d(x, y) = \max_{1 \leq i \leq n} |x_i - y_i|$

(iv) $d(x, y) = \sup_{1 \leq i \leq n} |x_i - y_i|$

where $x = (x_1, x_2, \dots, x_n)$

$y = (y_1, y_2, \dots, y_n)$

are any two points in \mathbb{R}^n .

(Choose the correct answer)

(j) Let (X, d_X) and (Y, d_Y) be two metric spaces and $f : X \rightarrow Y$ be continuous on X . Then for any $B \subseteq Y$.

(i) $f^{-1}(\overline{B}) \subseteq \overline{f^{-1}(B)}$

(ii) $\overline{f^{-1}(B)} \subseteq f^{-1}(\overline{B})$

(iii) $\overline{f(B)} \subseteq f(\overline{B})$

(iv) $f(\overline{B}) \subseteq \overline{f(B)}$

(Choose the correct answer)

2. Answer the following questions : $2 \times 5 = 10$

(a) Let $f(x) = x$ on $[0, 1]$ and

$$P = \left\{ x_i = \frac{i}{4}, i = 0, 1, \dots, 4 \right\}$$

Find $L(f, P)$ and $U(f, P)$.

(b) Let $f : [0, a] \rightarrow \mathbb{R}$ be given by

$f(x) = x^2$. Find

$$\int_0^a f(x) dx$$

- (c) Let (X, d) be a metric space and A, B be subsets of X . Prove that $(A \cap B)^0 = A^0 \cap B^0$.
- (d) If A is a subset of a metric space (X, d) , prove that $d(A) = d(\overline{A})$.
- (e) Let (X, d_X) and (Y, d_Y) be two metric spaces. Prove that if a mapping $f: X \rightarrow Y$ is continuous on X , then $f^{-1}(G)$ is open in X for all open subsets G of Y .

3. Answer **any four** parts : 5×4=20

(a) Prove that $f(x) = x^2$ on $[0, 1]$ is integrable.

(b) Show that $\lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{r}{r^2 + n^2} = \log \sqrt{2}$

(c) Let (X, d) be a metric space. Define $d': X \times X \rightarrow \mathbb{R}$ by

$$d'(x, y) = \frac{d(x, y)}{1 + d(x, y)} \text{ for all}$$

$x, y \in X$. Prove that d' is a metric on X .

(d) Let $X = C[a, b]$ and
 $d(f, g) = \sup\{|f(x) - g(x)| : a \leq x \leq b\}$
 be the associated metric where
 $f, g \in X$. Prove that (X, d) is a
 complete metric space.

(e) Let (X, d) be a metric space. Prove
 that a finite union of closed sets is
 closed.
 Infinite union of closed sets need not
 to be closed — Justify it. 3+2=5

(f) Let (X, d_X) and (Y, d_Y) be two metric
 spaces and $f : X \rightarrow Y$ be uniformly
 continuous. If $\{x_n\}_{n \geq 1}$ is a Cauchy
 sequence in X , prove that $\{f(x_n)\}_{n \geq 1}$
 is a Cauchy sequence in Y .

4. Answer **any four** parts : 10×4=40

(a) (i) Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous.
 Prove that f is integrable. 5

(ii) Discuss the convergence of the
 integral $\int_1^{\infty} \frac{1}{x^p} dx$ for various values
 at p . 5

- (b) (i) Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$. Prove that there exists $c \in [a, b]$ such that

$$\frac{1}{b-a} \int_a^b f(x) dx = f(c)$$

Using it prove that for $-1 < a < 0$ and $n \in \mathbb{N}$,

$$S_n = \int_a^0 \frac{x^n}{1+x} dx \rightarrow 0 \text{ as } n \rightarrow \infty$$

3+2=5

- (ii) Let $f : [a, b] \rightarrow \mathbb{R}$ be monotone. Prove that there exists $c \in [a, b]$ such that

$$\int_a^b f(x) dx = f(a)(c-a) + f(b)(b-c)$$

5

- (c) (i) Prove that a convergent sequence in a metric space is a Cauchy sequence.

Show that in the discrete metric space every Cauchy sequence is convergent.

3+2=5

- (ii) Define an open set in a metric space (X, d) .

Prove that in any metric space (X, d) , each open ball is an open set.

1+4=5

- (d) (i) Let (X, d) be a metric space and F be a subset of X . Prove that F is closed in X if and only if F^c is open in X . 5
- (ii) Let (X, d) be a metric space and Y a subspace of X . Let Z be a subset of Y . Prove that Z is open in Y if and only if there exists an open set $G \subseteq X$ such that $Z = G \cap Y$. 5
- (e) (i) Let (X, d_X) and (Y, d_Y) be metric spaces and $A \subseteq X$. Prove that a function $f : A \rightarrow Y$ is continuous at $a \in A$ if and only if whenever a sequence $\{x_n\}$ in A converges to a , the sequence $\{f(x_n)\}$ converges to $f(a)$. 6
- (ii) Prove that a mapping $f : X \rightarrow Y$ is continuous on X if and only if $f^{-1}(F)$ is closed in X for all closed subsets F of Y . 4

- (f) (i) Show that the function $f : (0, 1) \rightarrow \mathbb{R}$ defined by $f(x) = \frac{1}{x}$ is not uniformly continuous. 5
- (ii) Let (X, d) be a metric space and let $x \in X$ and $A \subseteq X$ be non-empty. Prove that $x \in A$ if and only if $d(x, A) = 0$. 5
- (g) (i) Define a connected set in a metric space. Prove that if Y is a connected set in a metric space (X, d) , then any set Z such that $Y \subseteq Z \subseteq \bar{Y}$ is connected. 1+4=5
- (ii) Let (X, d) be a metric space. Prove that the following statements are equivalent :
- (X, d) is disconnected
 - there exists a continuous mapping of (X, d) onto the discrete two element space (X_0, d_0) . 5

(h) Let (\mathbb{R}, d) be the space of real numbers with the usual metric. Prove that a subset I of \mathbb{R} is connected if and only if I is an interval.
