Total No. of printed pages = 6

3 (Sem 6) CHM M1

2015

32.75

CHEMISTRY

(Major)

Theory Paper: M-6.1

(Spectroscopy)

Full Marks - 60

Time – Three hours

The figures in the margin indicate full marks for the questions.

1. Answer the following questions:

 $1 \times 7 = 7$

- (a) What is the energy (in kJ mol⁻¹) of visible rays with $\lambda = 400$ nm?
- (b) Explain "H₂, N₂, O₂ etc molecules do not show pure rotational or vibrational spectra".
- (c) Which is not correct?
 - (i) Spherical top molecules have all three moments of inertia equal.

- (ii) Linear molecules have two moments of inertia equal and one is negligibly small.
- (iii) Asymmetric top molecules have two moments of inertia equal, but the other is not small.
- (d) What is a fundamental band in pure vibrational spectra?
- (e) What are Raman lines?
- (f) The absorbance of 5.4×10⁻⁴ M Fe³⁺ solution at 530 nm was 0.54. Calculate molar absorption coefficient, if the path length was 1 cm.
- (g) What is spin-spin relaxation in NMR spectroscopy?

2. Answer in short:

 $2\times4=8$

(a) The order of electronegativity in halogens is I < Br < Cl < F and proton NMR signals in CH, X(X = I, Cl, Br, F) are

> CH₃F CH₃Cl CH₃Br CH₃I 4.26δ 3.0δ 2.82δ 2.16δ

Explain the trend in NMR signals.

- (b) What are α-cleavage and induce cleavage in mass spectroscopy?
- (c) The moment of inertia of HC¹²N¹⁴ molecules is 1.89×10⁻⁴⁶ kgm². Calculate the rotational constant.
- (d) On what factors the relative intensities of spectral lines in rotational spectra are expected to depend on?
- 3. Answer any *three* questions of the following: $3\times5=15$
 - (a) Distinguish between emission and absorption spectra. State spectroscopic displacement law. Why the magnitude of energy of the electron in all the energy levels of an atom should decrease by a factor of 0.99945? 3+2=5
 - (b) What is rotational spectroscopy? How it differs from infrared spectroscopy? 4+1=5
 - (c) What is oscillator strength or f-number?

 Why it is necessary to introduce this number in electronic spectroscopy? How allowed and forbidden transitions are related to f-number?

 3+1+1=5

(d)	From the concept of quantum theory, prove
	that the frequency absorbed or emitted by a
	nucleus in moving from one energy level to
	another is directly proportional to the applied
	magnetic field. 5

- 4. Answer any *three* questions : $10 \times 3 = 30$
 - (a) (i) Explain with examples what are the types of ions produced in a mass spectrometer?
 - (ii) What is the use of 'Ring rule' in interpretation of mass spectra?
 - (b) (i) Discuss the theory of electron spin resonance spectroscopy. 7
 - (ii) Taking example of hydrogen atoms, explain what is meant by hyperfine splitting in electron spin resonance spectroscopy.
 - (c) (i) Define and explain what do you mean by equivalent hydrogens, coupling constant and up field and down field in NMR spectroscopy. 2+2+3=7
 - (ii) Predict how the high resolution ¹HNMR spectrum of each of following compounds will appear 3
 2-iodopropane and ethanol.

- (d) (i) "The absorption at longer wavelength in electronic spectroscopy is due to the combination of a chromophore and an auxochrome group." Describe what do you mean by chromophore and auxochrome. Also explain briefly red shift and blue shift.

 4+3=7
 - (ii) Predict λ_{max} of $\pi \rightarrow \pi^*$ transition for the following three molecules using Woodward's rule.

(e) (i) Explain how a non-linear N-atomic molecule and a linear N-atomic molecule can have 3N-6 and 3N-5 different internal vibrations respectively. Sketch the fundamental vibrations of water molecule, also show change in electric dipole moment produced by bending and symmetric vibrations of the molecule. 2+3+2=7

- (ii) The infrared spectrum of CO shows a vibrational absorption peak at 2170 cm⁻¹. What is the force constant for the CO bond.?
- (f) (i) Discuss the effect of isotopic substitution in rotational spectroscopy with respect to ¹²C ¹⁶O and ¹³C ¹⁶O molecules.

Draw and discuss Morse curve for a typical diatomic molecule. What is Morse function?

4+2+1=7

- (ii) Give spectral notation for the following states of the atom:

 3
 - (a) n=4, L=2, S=0
 - (b) n=4, L=1, S=1, J=0
 - (c) n=3, L=2 and multiplicity 2.